オレゴン州： 小麦作柄と気象状況
2010 年 7 月 25 日現在

概況　オレゴン州の 2010 年 7 月 25 日に終る 1 週間の気象は、高温・乾燥気象であった。Willamette Valley の最高気温は 92～96 度 F となり、平均気温は平年を 1～2 度 F 上回った。NC 地区の最高気温は 94～101 度 F となり、平均気温は平年を 1～2 度 F 上回った。1 週間の降水量は Portland にて 0.01 インチ記録されたが、他の観測所ではゼロであった。1 週間の州平均農作業稼働日数は 6.9 日（前週：6.9 日）であった。Topsoil 及び Subsoil の土壌水分は、前週より急激に減少し、“ Short ” と“ Very short が増えた。Willamette Valley の冬小麦は完熟し収穫直前となった。Sherman 郡及び Wasco 郡の北端では 7 月 19 日に冬小麦の収穫が開始された。Umatilla 郡の Pendleton の北、国道 11 号沿いで収穫が進んでおり、Athena 近辺（海抜 1500～1600 フィート）の冬小麦は高単収であると報告された。冬小麦は昨年の 2 週間遅れであり、全州の 20%が収穫された。春小麦は 97%が出穂し、概ね登熟期の後半となった。冬小麦の作柄は 72%が“ Good ”から“ Excellent ”であり、春小麦の作柄は 77%が“ Good ”から“ Excellent ”であった。

土壌水分(*)

(%)	Very short	Short	Adequate	Surplus
Topsoil	10	43	46	1
Last week	5	33	59	3
Subsoil	9	37	53	1
Last week	3	25	70	2

冬小麦の生育状況(*)

(%)	This week	Last week	Last year	5-Yr.Average
Harvested	20	8	49	42

冬小麦の作柄状況(*)

(%)	Very poor	Poor	Fair	Good	Excellent
This week	0	6	22	53	19
Last week	1	7	21	49	22

春小麦の生育状況(*)

(%)	This week	Last week	Last year	5-Yr.Average
Headed	97	94	100	99

春小麦の作柄状況(*)

(%)	Very poor	Poor	Fair	Good	Excellent
This week	1	4	18	45	32
Last week	1	5	17	41	36

*) Source: National Agricultural Statistic Service, USDA-Oregon Field Office. 5-yr average means average of 2005,2006,2007,2008 and 2009 crop years.

© Copyright 2002-2009 OMIC USA Inc. All rights reserved.
ご質問又はご意見は、OMIC USA Inc.小川正晃 ogawa.max@omicnet.comまでお願い致します。

2010年7月1日付けUSDA冬小麦生産量予想

State	Area Harvested 1,000 acres		Yield Bushels/acre			Production 1,000 bushels	
				2010			
	2009	2010	2009	June 1	July 1	2009	2010
Oregon	750	835	56.0	62.0	63.0	42,000	52,605
USA	34,485	32,085	44.2	46.6	46.9	1,522,718	1,505,493

2010年7月1日付けUSDA春小麦生産量予想

State	Area Harvested 1,000 acres		Yield Bushels/acre			Production 1,000 bushels	
				2010			
	2009	2010	2009	June 1	-	2009	2010
Oregon	127	130	54.0	55.0	-	6,858	7,150
USA	12,955	13,590	45.1	44.6	-	584,411	606,755

以上

© Copyright 2002-2009 OMIC USA Inc. All rights reserved.
ご質問又はご意見は、OMIC USA Inc.小川正晃 ogawa.max@omicnet.comまでお願い致します。